

Quaternary Fault and Fold Database of the United States

As of January 12, 2017, the USGS maintains a limited number of metadata fields that characterize the Quaternary faults and folds of the United States. For the most up-to-date information, please refer to the <u>interactive fault map</u>.

unnamed syncline west of Carbondale (Class B) No. 2335

Last Review Date: 1998-09-16

Compiled in cooperation with the Colorado Geological Survey

citation for this record: Widmann, B.L., compiler, 1998, Fault number 2335, unnamed syncline west of Carbondale, in Quaternary fault and fold database of the United States: U.S. Geological Survey website,

https://earthquakes.usgs.gov/hazards/qfaults, accessed 12/14/2020 03:00 PM.

Synopsis

This unnamed, east-northeast-trending synclinal sag lies within the Carbondale collapse center east of the Grand Hogback monocline. The sag is interpreted to result from flowage and dissolution of evaporite from beneath the area. Undivided Pleistocene "older " alluvium and colluvium are deformed by this structure (Kirkham and Widmann, 1997 #2711). There have not been any detailed studies conducted on this structure. In as much as the faulting may be aseismic (salt related), we considered this to be Class B structure.

County(s) and	Qf6 of Widmann and others (1998 #3441). JNTY, COLORADO
	CWALL COLD VIEW DAYS
Physiographic province(s) SOUTHERN RO	CKY MOUNTAINS
Reliability of location Good Compiled at 1:25	0,000 scale.
	synclinal sag was mapped at a scale of 1:24,000 Widmann (1997 #2711).
the Carbondale contained significant dissolution of evaluation and Widmann, 19	ast-northeast-trending synclinal sag lies within ollapse center, which is reported to have ificant Neogene collapse due to flowage and aporite deposits from beneath the area (Kirkham 1997#2711). The area is underlain by at least 900 eposits (Mallory, 1966 #2720).
Length (km) 1 km.	
Average strike N70°E	
Sense of movement Syncline Comments: Syncline the south and nor	line trends north-northeast; limbs dip roughly to th.
Dip Direction N; S	
Paleoseismology studies	
	synclinal sag developed in Pleistocene deposits idmann, 1997 #2711).

Age of faulted surficial deposits	Pleistocene "older" alluvium and colluvium are deformed by this synclinal sag (Kirkham and Widmann, 1997 #2711).				
Historic earthquake					
Most recent prehistoric deformation	undifferentiated Quaternary (<1.6 Ma) Comments: Pleistocene "older" alluvium and colluvium are deformed by this synclinal sag (Kirkham and Widmann, 1997 #2711).				
Recurrence interval					
Slip-rate category	Less than 0.2 mm/yr Comments: Widmann and others (1998 #3441) placed this anticline within the <0.2 mm/yr uplift-rate category.				
Date and Compiler(s)	1998 Beth L. Widmann, Colorado Geological Survey				
References	#2705 Kirkham, B., 1997, Late Tertiary and Quaternary collapse related to dissolution and flowage of Pennsylvanian evaporitic rocks in the Glenwood Springs area, Colorado, <i>in</i> McCalpin, J.P., ed., Active geologic environment of central Colorado, Aspen-Glenwood Springs-Silt, Colorado: Friends of the Pleistocene, Rocky Mountain Cell, September 12-14, 1997, Field guidebook. #2711 Kirkham, R.M., and Widmann, B.L., 1997, Geologic map of the Carbondale quadrangle, Garfield County, Colorado: Colorado Geological Survey Open-File Report 97-3. #2720 Mallory, W.W., 1966, Cattle Creek anticline, a salt diapir near Glenwood Springs, Colorado: U.S. Geological Survey Professional Paper 550-B, 12-15 p. #3441 Widmann, B.L., Kirkham, R.M., and Rogers, W.P., 1998, Preliminary Quaternary fault and fold map and database of Colorado: Colorado Geological Survey Open-File Report 98-8, 331 p., 1 pl., scale 1:500,000.				

Questions or comments?

Facebook Twitter Google Email

<u>Hazards</u>	_			
Design Ground M	<u> lotionsSeismic Hazaro</u>	l Maps & Site-S	pecific DataFa	ultsScenarios
	rdsDataEducationMor	•	•	
Search	Search			
HomeAbout UsCo	ontactsLegal			