

# Quaternary Fault and Fold Database of the United States

As of January 12, 2017, the USGS maintains a limited number of metadata fields that characterize the Quaternary faults and folds of the United States. For the most up-to-date information, please refer to the [interactive fault map](#).

## Buffalo Fork fault, southern section (Class A) No. 767c

Last Review Date: 1998-03-26

*citation for this record:* Pierce, K.L., and Machette, M.N., compilers, 1998, Fault number 767c, Buffalo Fork fault, southern section, in Quaternary fault and fold database of the United States: U.S. Geological Survey website, <https://earthquakes.usgs.gov/hazards/qfaults>, accessed 12/14/2020 02:02 PM.

### Synopsis

**General:** The Laramide Buffalo Fork thrust fault is an important crustal feature that thrusts upper Paleozoic sedimentary rocks westward onto Cretaceous rocks. Parts of this thrust have been reactivated in Cenozoic time with down-to-the-east normal faulting, including one site of observed post-glacial fault scarps and another of tilted late Quaternary lake sediment. This fault is in a remote area seldom visited by geologists that are experienced in neotectonics; thus, evidence of Quaternary movement may have been missed. Because of its older history, this structure might serve to accommodate ongoing tectonic activity. There have been several Quaternary basalt extrusions in the hanging wall of the fault.

**Sections:** This fault has 3 sections. Fault divided on basis of young activity in the middle section [767b] and older movement

	on the ends [northern section, 767a; southern section, 767c].
<b>Name comments</b>	<p><b>General:</b> Referred to as the Buffalo Fork thrust (fault) by Love and Keefer (1975 #2285). The Buffalo Fork thrust bounds the east margin of the Two Oceans Plateau and extends from Yellowstone Lake on the north to Togowotee Pass area on the south. The northern part [767a] represents normal-fault reactivation of Buffalo Fork thrust fault. Although this fault was also referred to as the South Arm fault by Wong and others (2000 #4484), the Buffalo Fork fault name has precedence and is used herein.</p> <p><b>Section:</b> This informally named section of the Buffalo Fork fault is a southern extension of the fault beyond the middle section that probably has post-glacial scarps. Shown by Love and Christiansen (1985 #2287) and Christiansen (2001 #1784) as having normal fault movement for 18 km south of the Yellowstone National Park boundary. This informally named section of the Buffalo Fork fault is a southern extension of the fault beyond the middle section that probably has post-glacial scarps. Shown by Love and Christiansen (1985 #2287) and Christiansen (2001 #1784) as having normal fault movement for 18 km south of the Yellowstone National Park boundary.</p> <p><b>Fault ID:</b> Refers to fault 237 (unnamed) of Witkind (1975 #819).</p>
<b>County(s) and State(s)</b>	TETON COUNTY, WYOMING
<b>Physiographic province(s)</b>	MIDDLE ROCKY MOUNTAINS
<b>Reliability of location</b>	<p>Poor Compiled at 1:125,000 scale.</p> <p><i>Comments:</i> Mapped by Love and Christiansen (1985 #2287) at 1:500,000 scale and northernmost part of section is shown by Christiansen (2001 #1784) at 1:125,000 scale. Fault traces recompiled at 1:125,000 scale on map with topographic base. Trace is good for the older thrust fault, but uncertain (poor) for the normal fault.</p>
<b>Geologic setting</b>	The Buffalo Fork thrust is a major Laramide structure in northwest Wyoming (Love and Keefer, 1975 #2285). Late Cenozoic extension may have been accommodated along or near this zone of structural weakness (p. D45-46 in Love and Keefer, 1975 #2285). Christiansen (2001 #1784) has recently affirmed

	<p>this reactivation. Because of its older history, this structure might serve to accommodate ongoing tectonic activity. There have been several Quaternary basalt extrusions in the hanging wall of the fault (U.S. Geological Survey, 1972 #639; Christiansen, 2001 #1784).</p>
<b>Length (km)</b>	This section is 2 km of a total fault length of 30 km.
<b>Average strike</b>	N9°W (for section) versus N1°E (for whole fault)
<b>Sense of movement</b>	<p>Normal</p> <p><i>Comments:</i> Love and Keefer (1975 #2285) and Christiansen (plate 3, 2001 #1784) show this as a Laramide thrust fault with later (Quaternary) normal fault movement.</p>
<b>Dip Direction</b>	<p>E</p> <p><i>Comments:</i> No known scarps on surficial material.</p>
<b>Paleoseismology studies</b>	
<b>Geomorphic expression</b>	
<b>Age of faulted surficial deposits</b>	Observed mostly in pre-Quaternary bedrock as thrust with subsequent late Cenozoic ? reactivation as a normal fault.
<b>Historic earthquake</b>	
<b>Most recent prehistoric deformation</b>	<p>undifferentiated Quaternary (&lt;1.6 Ma)</p> <p><i>Comments:</i> No Quaternary offset has been established, but section included herein based on association with young movement on adjacent (middle) section of Buffalo Fork fault.</p>
<b>Recurrence interval</b>	
<b>Slip-rate category</b>	<p>Less than 0.2 mm/yr</p> <p><i>Comments:</i> Wong and others (2000 #4484) suggested fault slip rates ranging from 0.4-1.4 mm/yr, each with separate weighting. However, these rates are based on an assumption of activity</p>

	<p>identical to their appraisal of the Yellowstone Lake (Eagle Bay [757]) fault. Wong and others' (2000 #4484) reported slip rates are model dependent and do not represent actual measured values. The late Quaternary characteristics of this fault (overall geomorphic expression, lack of scarps on Quaternary materials, age of faulted deposits, etc.) suggest the slip rate during this period is of a lesser magnitude. Accordingly, the &lt;0.2 mm/yr slip-rate category has been assigned to this fault.</p>
<p><b>Date and Compiler(s)</b></p>	<p>1998  Kenneth L. Pierce, U.S. Geological Survey, Emeritus  Michael N. Machette, U.S. Geological Survey, Retired</p>
<p><b>References</b></p>	<p>#1784 Christiansen, R.L., 2001, The Quaternary and Pliocene Yellowstone Plateau volcanic field of Wyoming, Idaho, and Montana: U.S. Geological Survey Professional Paper 729-G, 145 p., 3 pls., scale 1:125,000.</p> <p>#2287 Love, J.D., and Christiansen, A.C., 1985, Geologic map of Wyoming: State Geologic Map, 3 sheets, scale 1:500,000.</p> <p>#2285 Love, J.D., and Keefer, W.R., 1975, Geology of sedimentary rocks in southern Yellowstone National Park, Wyoming: U.S. Geological Survey Professional Paper 729-D, 60 p.</p> <p>#639 U.S. Geological Survey, 1972, Geologic map of Yellowstone National Park: U.S. Geological Survey Miscellaneous Geologic Investigations I-711, 1 sheet, scale 1:125,000.</p> <p>#819 Witkind, I.J., 1975, Preliminary map showing known and suspected active faults in Wyoming: U.S. Geological Survey Open-File Report 75-279, 35 p. pamphlet, 1 sheet, scale 1:500,000.</p> <p>#4484 Wong, I., Olig, S., and Dober, M., 2000, Preliminary probabilistic seismic hazard analyses—Island Park, Grassy Lake, Jackson Lake, Palisades, and Ririe Dams: U.S. Department of the Interior, Bureau of Reclamation Technical Memorandum D8330-2000-17.</p>

[Questions or comments?](#)

[Facebook](#) [Twitter](#) [Google](#) [Email](#)

[Hazards](#)

[Design Ground Motions](#)[Seismic Hazard Maps & Site-Specific Data](#)[Faults](#)[Scenarios](#)

[Earthquakes](#)[Hazards](#)[Data](#)[Education](#)[Monitoring](#)[Research](#)

[Home](#)[About Us](#)[Contacts](#)[Legal](#)