



**SURFICIAL MATERIAL GEOLOGIC MAP OF THE WEBSTER GROVES 7.5' QUADRANGLE
ST. LOUIS CITY AND COUNTY, MISSOURI AND ILLINOIS**

Geology and Digital Compilation by
David A. Gaunt and Travis Carr

2010

OFM-10-562-GS



MISSOURI DEPARTMENT OF NATURAL RESOURCES
DIVISION OF GEOLOGY AND LAND SURVEY
GEOLOGICAL SURVEY PROGRAM
P.O. BOX 250, ROLLA MO 65402-0250
www.dnr.mo.gov/geology
573-368-2100

**THIS MAP WAS PRODUCED UNDER A COOPERATIVE
AGREEMENT WITH THE UNITED STATES GEOLOGICAL
SURVEY**



Permission must be obtained to visit privately owned land

PHYSIOGRAPHY

The Webster Groves quadrangle includes part of the floodplain of the Mississippi River and loess covered uplands. The floodplain is less than one mile wide in this area with only a small portion residing on this quadrangle. The quadrangle lies within the Dissected Till Plains Section of the Central Lowland Province of the Interior Plains Physiographic Division. The lowest recorded elevation of slightly less than 380 feet mean sea level (msl) occurs along the edge of the Mississippi River. The highest elevation on the quadrangle occurs on the loess covered uplands and is slightly greater than 653 feet msl in the eastern portion of the quadrangle. Total relief on the Webster Groves quadrangle is approximately 273 feet.

DESCRIPTION OF MAP UNITS


- AF** **ARTIFICIAL FILL** – This unit comprises artificially emplaced fill material and is composed of a mixture of heterogeneous clay, silt, sand and gravel in various quantities. This unit may reach 40 feet in total thickness and comprises the material for highway and railroad beds and waste water treatment facility fill. This artificial fill has typically been placed on undisturbed materials.
 - Qcly** **QUATERNARY CLAY-CAPPED ALLUVIUM** – This unit has been deposited by the Mississippi River. The approximate upper 15 feet of these deposits are composed predominantly of clay with variable amounts of silt and organic material. The material residing below the clay is predominantly sand and gravel to the top of bedrock and can be up to 130 feet thick near the large river. The water table is approximately five to 15 feet below ground surface here resulting in an interval of saturated sand and gravel greater than 100 feet thick.
 - Qsilt** **QUATERNARY SILT-CAPPED ALLUVIUM** – This unit has been deposited by the Mississippi River and its tributaries. The approximate upper 15 feet of these deposits are composed predominantly of silt with variable amounts of clay and organic material. The material residing below the silt is predominantly sand and gravel to the top of bedrock and can reach 130 feet thick. The water table is approximately five to 15 feet below ground surface resulting in an interval of saturated sand and gravel greater than 100 feet thick.
 - Qsnd** **QUATERNARY SAND-CAPPED ALLUVIUM** – This unit has been deposited by the Mississippi River and its tributaries. The composition of this unit is predominantly sand with variable amounts of clay, silt and organic material in the upper 15 feet. In the eastern portion of the map in Illinois. The thickness of this unit reaches 130 feet along the large rivers. The water table is approximately 15 feet below ground surface, resulting in an interval of saturated sand greater than 100 feet thick.
 - Ql** **QUATERNARY LOESS** – This unit is a wind-blown deposit of silt and clayey silt with occasional pockets of clay, sand and gravel. The unit is composed of two separate loess layers, the Roxana below and the Peoria above (Goodfield, 1965). The total thickness of the two units may reach 70 feet. The Roxana is higher in clay content and may have a paleosol developed in the upper few feet. The contact between the two units forms a potential slide plane in areas of higher slope. The loess overlies Mississippian-age bedrock. The Mississippian units are limestone and shale creating unique environments. Where the loess is thin, the limestone may be karstic. Where the underlying unit is predominantly shale, water will perch destabilizing the contact zone. Where the loess rests upon shale, the slide potential is increased.
 - Qt** **QUATERNARY TILL** – Deposits of clayey till are located in the western portion of the quadrangle. The quaternary till was deposited as a drift blanket during glaciation north of the Missouri River. The till is a mixture of clay, silt, sand, gravel and cobbles that covers the bedrock surface. The till varies in thickness from 10 to 25 feet with the thickest deposits inland from the river and in depressions of the bedrock surface.
 - Qtd** **QUATERNARY TERRACE DEPOSIT** – The terrace deposits in the quadrangle are slightly different than previously mapped terrace deposits. All were deposited during glacial events leaving the terrace above low flow stage of the river. However, the terrace deposits in this quadrangle have a lacustrine signature of sensitive organic clays approximately 20 feet below the surface. After high stage flow dropped to normal, low lying areas within the terrace were filled with organic clay material. This zone has a very low shear wave velocity and lies under infrastructure in many areas.
 - K** **KARST** – These areas have high concentrations of sinkholes, caves and other karstic features due to the solutional weathering of the Mississippian-age limestone underlying this area. These areas are typically found in the upland regions of the quadrangle.
 - B** **BEDROCK** – The exposures in this quadrangle are typically found in areas of high topographic relief, near tributaries or in quarries.
 - R** **RESIDUUM** – The areas of residual material are found on the slope and toe slope of high relief areas. The content is dependent on the bedrock parent material but typically has a high clay content with fragments of parent material.
- A—A'** Line locates the placement of the cross section with end line symbols.




BIBLIOGRAPHY

- Allen, W.H. and Ward, R.A., 1977, Soil, in The Resources of St. Charles County, Missouri, land, water, and minerals; Satterfield, Ira and Harris, Barbara, eds.; Missouri Geological Survey, Department of Natural Resources, 237 p.
- Goodfield, A.G., 1965, Pleistocene and surficial geology of the City of St. Louis and the adjacent St. Louis County, Missouri; unpublished Ph.D. dissertation, University of Illinois, Urbana, IL, 206p., 6 pl.
- Illinois State Geological Survey, Water and related wells in Illinois, IGS map service: ILWATER 5/23/2007 <http://ablation.isgs.uiuc.edu/webSite/water/viewer.htm>
- Missouri Department of Natural Resources, 2007, Well Logs, Wells Certified, Bedrock, Roads, IMOP, in Missouri Environmental Geology Atlas (MEGA), Division of Geology and Land Survey, Missouri Department of Natural Resources.
- Schrader, W.D., and Krusekoph, H.H., 1956, Soil survey of St. Charles County, Missouri; Soil Conservation Service, U.S. Department of Agriculture, 49 pages, 1 pl.
- Thompson, Thomas L., 1995, The stratigraphic succession in Missouri, v. 40 rev.; Division of Geology and Land Survey, Missouri Department of Natural Resources, 190 p.
- Williams, R.A., Odum, J.K., Stephenson, W.J., and Herrmann, R.B., 2007, Shallow P- and S-wave velocities in the St. Louis region, Missouri-Illinois; Earthquake Spectra, v. 23, no. 3, 711-726.

SURFICIAL MATERIAL THICKNESS



TOP OF BEDROCK



DATA POINT LOCATION

