



# Comparisons of Darfield and Christchurch Ground Motions with NGA-W1 GMPEs

David M. Boore



(<http://www.geonet.org.nz/canterbury-quakes/>)


## Stations for which $R_{jb} \leq 200$ km



Class C sites at greater distance may have smaller depth of sediments than those near Christchurch

Most class C sites to the north

- Many sites with  $V_{S30} = 275$  and  $660$  m/s because values were estimated, not measured
- Distributions of  $V_{S30}$  similar for both events
- Most records with  $R < 100$  km are from class D sites



Note: Some sites classified as “E” by Bradley & Cuprinovski (2011)


# Comparison of Darfield and Christchurch Ground Motions

- Motions from both events at close distances are comparable
- Apparently no or small site effect (but note difference in spatial locations for different class sites, so source effects could compensate for site effects)
- Nonlinear soil response could have reduced motions for the larger event


Observed RotD50 reduced to GMRotI50 using factors from Boore (2010); maximum effect is 4% at T=5 s.

In this and subsequent comparisons, the Y-axis spans three orders of magnitude.


Concentrate on H components, as the 2008 NGA GMPEs were only for H.



- Christchurch (**M** 6.1) class D motions comparable or even larger than Darfield (**M** 7.0) at close distances
- Darfield class C motions greater than Christchurch motions ( $R > 50$  km)




- Christchurch (**M** 6.1) class D motions now smaller than Darfield (**M** 7.0) at all distances
- Apparent site effect small or not existent






# Comparison of Observed Motions and Motions from NGA-W1 GMPEs: Darfield


- Reddish GMPEs for class D
- Bluish GMPEs for class C
- Don't try to follow curves for individual GMPEs
- Overall comparison good
- Apparent lack of site effect in data is consistent with GMPEs



- Note separation in observed class D values; GMPEs predictions between the two groups
- Site effect in GMPEs, but not apparent in observations (but little overlap in distance range for class C and D)




- GMPEs underpredict observations (except class C at greater distances)
- Site effect in GMPEs, but not apparent in observations (but little overlap in distance range for class C and D)




Comparison of Observed  
Motions and Motions from  
NGA-W1 GMPEs:  
Christchurch


- Overall comparison good
- Apparent lack of site effect in data is consistent with GMPEs



- GMPEs tend to underpredict class D motions at close distances
- Site effect in GMPEs, but not apparent in observations (but little overlap in distance range for class C and D)




- GMPEs severely underpredict observations at shorter distances
- Agreement better for greater distances
- Site effect in GMPEs, but not apparent in observations (but little overlap in distance range for class C and D)






# Sensitivity of Predicted $T=5$ s PSA to sediment depth and to magnitude


- Show AS08 and class D only
- $Z_{1.0}=1000$  m from Bradley & Cuprinovski (2011)
- **M** 6.33 from Holden (2011)
- Use of these data in NGA-W2 without  $Z_{1.0}$  and larger **M** could result in biased results.
- Does  $Z_{1.0}$  vary spatially, with it being smaller for stations at greater distances? If so, this would help explain the discrepancy at greater distance. This could also be due to a difference in geometrical spreading due to lateral changes in crustal structure.



# Vertical Motion



(Fry et al., 2011)




(Bradley & Cuprinovski, 2011)

# Effects Producing Spatial Variability in Ground Motions

- Source: Radiation Pattern & Directivity
- Path: volcanic vs sediments
- Basin Waves
- Sediment Depth
- Shallow Site Response
  - Linear
  - Nonlinear


# Fault Normal and Fault Parallel Velocity Time




- Drastic change in frequency content going from sites on rock (HVSC & LPCC) to sites on Quaternary sediments.

# Evidence for Nonlinear Soil Response






### 2001 M 6.8 Nisqually (SDS)



(see Frankel et al., 2002, for details)



Negative vertical accelerations are “clipped”. This may be due to a different nonlinear process than that producing the cusps shown in the previous figures.


(Fry et al., 2011)

# Conclusions



- M 7.0 Darfield and M 6.1 Christchurch motions similar for close distances, short periods
- M 7.0 Darfield motions higher than M 6.1 Christchurch motions for longer periods (as expected from the difference in magnitudes)
- Site response not too obvious, but this may be because of the different spatial distributions of the site classes (most close sites are class D)
- Observed motions influenced by many effects, including
  - Lateral changes in geology
  - Local linear and nonlinear site response
  - Basin waves (?)
- GMPEs are in reasonable agreement with observations for close distances, short periods
- GMPEs underpredict longer period motions, using metadata in current NGA-W2 flatfile

**END**






From B. Chiou, Source: GNS Science




(Bradley & Cuprinovski, 2011)






(Bradley & Cuprinovski, 2011)

# Directivity Effect and Velocity Pulse



Source: George Walker

(from B. Chiou)




(Bradley & Cuprinovski, 2011)



(Bradley & Cuprinovski, 2011)

# 0282 Christchurch, New Zealand



(from B. Chiou)


# Fault Rupture


- Reverse faulting on a buried fault
- Assumed fault plane
  - Strike =  $68^\circ$  (from USGS CMT)
  - $65^\circ$  dip, to the south
  - Top of rupture is at 2 km depth (assumed)
  - Bottom of rupture is at 12 km (assumed)
  - Rupture length  $\sim$  15 km (length of the aftershock zone).

(from B. Chiou)


## 2001 Nisqually, Washington, earthquake (M 6.8)


- SDS within 200 m of SDN
- liquefaction at SDS, not at SDN
- Note cusps at SDS and increased amplitude at high frequencies















(Bradley & Cuprinovski, 2011)

Stations for  
which  $R_{jb} \leq$   
200 km











Observed RotD50 reduced to GMRot150 using factors from Boore (2010); maximum effect is 4% at T=5 s.


In this and subsequent comparisons, the Y-axis spans three orders of magnitude.


Concentrate on H components, as the 2008 NGA GMPEs were only for H.




















(Fry et al., 2011)